

If you have an Injectafer prescription, you may be able to get help with your out-of-pocket costs.

Injectafer is the **only** IV iron with a copay assistance program.*

THE INJECTAFER SAVINGS PROGRAM

If you're eligible,† you may:

- Receive assistance of up to \$500 per dose with a maximum benefit of up to \$1000 per course of treatment (2 doses)
- Stay enrolled for 2 courses of treatment (4 doses) per 12-month period

For each course of treatment:

Get your
FIRST DOSE
for as little as
\$50
Up to 750 mg of iron
for qualified patients†

Get your
SECOND DOSE
for as little as
\$0
Up to 750 mg of iron
for qualified patients†

Doses are separated by at least 7 days.
Note: The first dose of the second course may cost as little as \$50.



Have questions about the Injectafer Savings Program?

Call **1-877-4-IV-IRON (1-877-448-4766)**

Available Monday–Friday, 8:00 AM–5:00 PM ET

*As of October 2019.

†The Injectafer Savings Program is only available for adults 18 years or older who are commercially insured or cash-paying patients. Insurance out of pocket payment must be over \$50. Other restrictions may apply. Please see full Terms and Conditions.

‡If you weigh less than 50 kg (110 lb), your doctor will prescribe a lower dose of Injectafer for you.

INDICATIONS

Injectafer® (ferric carboxymaltose injection) is an iron replacement product indicated for the treatment of iron deficiency anemia (IDA) in adult patients who have intolerance to oral iron or have had unsatisfactory response to oral iron, and in adult patients with non-dialysis dependent chronic kidney disease.

SELECTED SAFETY INFORMATION ABOUT INJECTAFER

CONTRAINDICATIONS

Injectafer is contraindicated in patients with hypersensitivity to Injectafer or any of its inactive components.

Please [click here](#) for full Prescribing Information.



ARE YOU ELIGIBLE?

- ✓ You have commercial insurance OR ✓ You pay for treatment with cash
- ✗ You have Medicare, Medicaid, or other federal or state healthcare insurance OR ✗ You have private indemnity or HMO insurance that reimburses patients for the entire cost of prescription drugs OR ✗ You are Medicare-eligible and enrolled in an employer-sponsored health plan

To see if you're eligible, visit www.injectafercopay.com or call **1-877-4-IV-IRON** (1-877-448-4766).

HOW TO GET YOUR SAVINGS

- 1** Ask your doctor to enroll you. They will need the last 4 digits of your Social Security number to confirm you're eligible.
If your doctor cannot enroll you, you can enroll yourself in one of two ways:
 - Visit injectafercopay.com OR • Call the IV Iron Hotline (1-877-448-4766)
 - ★ The best way for you to enroll
- 2** Once you're enrolled and receive your infusion of Injectafer, your doctor will bill the Injectafer Savings Program for your copay requirement.

If the copay requirement is more than \$500 for your infusion, your doctor may bill you for the remaining amount that you'll need to pay.

If your doctor requires you to pay the copay when you receive your infusion, the Injectafer Savings Program will reimburse you via paper check. To do this, you will need a Check Request Form. The best place to get the form is online at injectafercopay.com. If you have any questions throughout the process, please call the IV Iron Hotline (1-877-448-4766).

Injectafer Savings Program contact information

Phone: 1-877-4-IV-IRON (1-877-448-4766)

Fax: 1-888-257-4673

Website: www.injectafercopay.com

Mail: Injectafer Savings Program
100 Passaic Ave, Suite 245
Fairfield, NJ 07004

SELECTED SAFETY INFORMATION ABOUT INJECTAFER
WARNINGS AND PRECAUTIONS

Serious hypersensitivity reactions, including anaphylactic-type reactions, some of which have been life-threatening and fatal, have been reported in patients receiving Injectafer.

Please [click here](#) for full Prescribing Information.



Terms and conditions

1. This offer is valid for commercially-insured as well as cash paying patients.
2. Depending on insurance coverage, eligible insured patients may pay no more than \$50 for Injectafer for the first dose and \$0 for Injectafer for the second dose, up to a maximum savings limit of \$500 per dose, a \$1,000 program limit per course of therapy. Check with your pharmacist or healthcare provider for your copay discount. Patient out-of-pocket expense may vary.
3. This offer is not valid for patients enrolled in Medicare, Medicaid, or other federal or state healthcare programs, or private indemnity or HMO insurance plans that reimburse you for the entire cost of your prescription drugs. Patients may not use this card if they are Medicare-eligible and enrolled in an employer-sponsored health plan or medical or prescription drug benefit program for retirees.
4. The offer is valid for 2 courses, or 4 doses, of an Injectafer prescription. An explanation of benefits statement must be faxed in prior to transacting on the account numbers for assistance. One enrollment is allowed per 12-month period.
5. Daiichi Sankyo, Inc. reserves the right to rescind, revoke, or amend this offer without notice.
6. Offer good only in the USA, including Puerto Rico, at participating pharmacies or healthcare providers.
7. Void if prohibited by law, taxed, or restricted.
8. This account number is not transferable. The selling, purchasing, trading, or counterfeiting of this account number is prohibited by law.
9. This account number is not insurance.
10. By redeeming this account number, you acknowledge that you are an eligible patient and that you understand and agree to comply with the terms and conditions of this offer.
11. Qualified patients receiving Injectafer will be allowed a 60-day retroactive enrollment period to receive benefits under the program rules.

SELECTED SAFETY INFORMATION ABOUT INJECTAFER

WARNINGS AND PRECAUTIONS (cont'd)

Patients may present with shock, clinically significant hypotension, loss of consciousness, and/or collapse. Monitor patients for signs and symptoms of hypersensitivity during and after Injectafer administration for at least 30 minutes and until clinically stable following completion of the infusion. Only administer Injectafer when personnel and therapies are immediately available for the treatment of serious hypersensitivity reactions.

Please [click here](#) for full Prescribing Information.



Injectafer® (ferric carboxymaltose injection) is available by prescription only. Ask your doctor or healthcare provider if Injectafer is right for you.

What is Injectafer?

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

It is not known if Injectafer is safe and effective for use in children.

IMPORTANT SAFETY INFORMATION

Who should not receive Injectafer?

You should not receive Injectafer if you are allergic to ferric carboxymaltose or any of the other ingredients in Injectafer. The active ingredient in Injectafer is ferric carboxymaltose, the inactive ingredients are: water for injection, sodium hydroxide and/or hydrochloric acid.

What should I tell my doctor or healthcare provider before receiving Injectafer?

Before you receive Injectafer, tell your healthcare provider about all of your medical conditions, including if you:

- Have had an allergic reaction to iron given intravenously (into your vein), including Injectafer, or to other non-oral iron treatments
- If you have, or have previously experienced, iron overload, or if your body has difficulty using iron appropriately
- Have high blood pressure
- Are pregnant or plan to become pregnant. It is not known if Injectafer will harm your unborn baby. Your healthcare provider will decide if it is safe for you to take Injectafer
- Are breastfeeding or plan to breast feed. Injectafer passes into your breast milk. It is unknown whether Injectafer would pose a risk to your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with Injectafer.

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of Injectafer?

Injectafer can cause serious side effects, including:

- **Serious allergic reactions** that may be life-threatening, including shock, low blood pressure, loss of consciousness, and death.

Your doctor or healthcare provider will monitor you for signs and symptoms of an allergic reaction during and after each dose of Injectafer for at least 30 minutes. Other serious allergic reactions include itching, rash, hives, wheezing, or low blood pressure. You should report any signs and symptoms of an allergic reaction to Injectafer, in particular rashes, shortness of breath and wheezing to your doctor or healthcare provider.

- **High blood pressure**, sometimes with facial flushing, dizziness, or nausea, has been seen during treatment with Injectafer. This increase in blood pressure typically resolves within 30 minutes. Your doctor or healthcare provider will monitor you for signs and symptoms of an increase in blood pressure following each use of Injectafer.

Other serious side effects that have been reported include rash, difficulty breathing, itching, rapid heartbeat, fever, chest discomfort, chills, swelling of the face, lips, or tongue, back pain, muscle aches, and fainting.

The most common side effects of Injectafer include:

- Nausea, high blood pressure, flushing, low levels of phosphorous in your blood, dizziness, vomiting, headache, an increase in certain liver enzymes, and pain or bruising at the injection site. Potentially long-lasting brown staining of skin near the injection site may occur if Injectafer leaks out of the vein.

Excessive amounts of Injectafer may lead to a condition called iron overload, which is a buildup of iron and may be harmful.

These are not all of the possible side effects of Injectafer.

Tell your doctor if you have any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects.

General information about Injectafer

Injectafer may impact laboratory tests that measure iron in your blood for 24 hours after receiving Injectafer. Let your healthcare provider and laboratory staff know if you have received Injectafer within 24 hours of having blood tests.

To report side effects, contact American Regent at 1-800-734-9236 or E-mail: pv@americanregent.com or Fax: 1-610-650-0170.

You may also report side effects to the FDA at 1-800-332-1088 or www.fda.gov/medwatch.

The risk information provided here is not comprehensive. To learn more about Injectafer, talk with your healthcare provider or pharmacist. The FDA-approved product labeling can be found at www.injectafer.com/pdf/pi.pdf or call 1-800-645-1706.

Please see [Full Prescribing Information](#) for Injectafer.



Daiichi-Sankyo

